arXiv:1109.6275v1 [math.CO] 28 Sep 2011

A CLASSIFICATION OF ALL 1-SALEM GRAPHS

JONATHAN COOLEY, LEE GUMBRELL, AND JAMES MCKEE

Abstract. A cyclotomic graph is one that has all its eigenvalues in the interval $[-2,2]$, and a Salem graph is either (i) bipartite with all but two eigenvalues in $[-2,2]$, or (ii) non-bipartite with all but one eigenvalue in the interval $[-2,2]$. We define an m-Salem graph to be a connected Salem graph G for which m is minimal such that there exists an induced cyclotomic subgraph of G that has m fewer vertices than G. Every Salem graph contains a 1-Salem graph as an induced subgraph. The main result of this paper is a complete combinatorial description of all 1-Salem graphs: there are 25 infinite families and 383 sporadic examples, only 16 of which give trivial Salem graphs. We observe that all generalised line graphs that are Salem graphs contain at least one triangle, and hence produce a complete list of all 25 connected non-bipartite triangle-free Salem graphs.

1. Introduction

The notion of a *graph Salem number* was introduced by McKee and Smyth in [13]: these are Salem numbers associated to certain *Salem graphs* (precise definitions are recalled in Section 2). A number of partial results concerning the structure of Salem graphs appear in [13, Section 3], and a complete description of all Salem trees in [13, Theorem 7.2], but a combinatorial description of all Salem graphs is still unknown. In this paper we make fresh progress towards this goal; we partition the set of all Salem graphs into the *m-Salem graphs* (Definition 1) for $m = 1, 2, 3, \ldots$, and give a complete description of the first of these subsets (the 1-Salem graphs: Theorem 1). Part of the significance of this is that every Salem graph contains a 1-Salem graph as an induced subgraph, so Theorem 1 provides a stronger description of some necessary substructure of a Salem graph than had previously been known.

A graph is called *cyclotomic* if all its eigenvalues lie in the interval $[-2,2]$. A complete description of all cyclotomic graphs was provided by Smith in [17]. Every graph contains a cyclotomic graph as an induced subgraph, and it is a natural question to ask what is the smallest number of vertices that need to be deleted in order to produce such a graph. With Salem graphs in mind, this motivates the following definition.

Definition 1. *A connected Salem graph G is called an m-*Salem graph *if m is minimal such that there exists a set of m vertices $v_1, \ldots, v_m$ for which the induced graph $G - \{v_1, \ldots, v_m\}$ is cyclotomic.*

We shall need to consider bipartite graphs and non-bipartite graphs separately, as the definition of a Salem graph differs between the two cases (Definition 2, below). For non-bipartite Salem graphs all eigenvalues are at least -2, and Cameron *et al* ([2]) showed that the connected graphs with this property are either generalized

---

1991 *Mathematics Subject Classification.* 11R06, 05C50.

line graphs or belong to a finite list of graphs represented in the root system E_8. We summarise our main results in the following Theorem.

Theorem 1. *(i) All bipartite* 1*-Salem graphs are described by Theorem 11.*

(ii) Every 1*-Salem graph that is a generalised line graph is described by Theorem 12; there are* 25 *infinite families and* 6 *sporadic examples.*

(iii) There are 377 *non-bipartite* 1*-Salem graphs that are not generalised line graphs: see Theorem 18.*

In Section 2 we state some definitions and results that we will need to prove the three components of Theorem 1. In Section 3 we show that there exist non-trivial m-Salem graphs for any $m \geq 1$. The proof of Theorem 1 is split over three sections: Section 4 for the bipartite graphs; Section 5 for the generalised line graphs; and Section 6 for the non-bipartite graphs that are not generalised line graphs. The proofs for the latter two are split into further subsections. Observations on triangle-free non-bipartite Salem graphs and trivial Salem graphs are made in Sections 7 and 8, respectively.

2. BACKGROUND

We follow [13] in our definition of a *Salem graph*, splitting into two cases according as the graph is bipartite or not. Most of our interest will be in the non-bipartite case, with bipartite graphs constrained to the brief Section 4. The set of eigenvalues of a graph is simply the union of those coming from its connected components, so we shall mainly be considering connected graphs, but when we consider deleting a vertex the resulting induced subgraph might be disconnected, and consequently it is convenient to allow disconnected graphs in Definition 2, as well as in our definition of cyclotomic graphs. The assertions within Definition 2 concerning Salem numbers are established in [13].

Definition 2. *A bipartite graph G is called a* Salem graph *if the largest eigenvalue (or* index*) λ_1 is greater than* 2*, and the remaining $n-1$ eigenvalues are ≤ 2. Then $\lambda_n = -\lambda_1 < -2$. A bipartite Salem graph is called* trivial *if $\lambda_1^2 \in \mathbb{Z}$. The associated number $\tau(G)$ is the larger root of $\sqrt{z} + 1/\sqrt{z} = \lambda_1$; this is a Salem number unless G is trivial (in which case it is a quadratic Pisot number).*

A non-bipartite graph G is called a Salem graph *if the largest eigenvalue $\lambda_1 > 2$ and the remaining $n-1$ eigenvalues are in the interval $[-2, 2]$. A non-bipartite Salem graph is called* trivial *if $\lambda_1 \in \mathbb{Z}$. The associated number $\tau(G)$ is the larger root of $z + 1/z = \lambda_1$; this is a Salem number unless G is trivial.*

We recall Smith's classification of cyclotomic graphs.

Lemma 2 (Smith [17])**.** *The connected cyclotomic graphs are precisely the induced subgraphs of the graphs $\tilde{E}_6$, $\tilde{E}_7$, $\tilde{E}_8$, $\tilde{A}_n$ $(n \geq 2)$ and $\tilde{D}_n$ $(n \geq 4)$, shown in Figure 1.*

We refer to the following result as "interlacing". It allows us to bound the second-largest eigenvalue of a graph by the index of any induced subgraph obtained by deleting a single vertex.

Theorem 3 (Cauchy [4]; or see [9], Theorem 9.1.1)**.** *Let G be an n-vertex graph with vertex set $V(G)$ and eigenvalues $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_n$. Also, let H be the induced*



FIGURE 1. The maximal connected cyclotomic graphs. Note that each graph has one more vertex than the subscript in its name.

graph on $V(G) \setminus \{v\}$ *obtained from* G *by deleting the vertex* v *and its incident edges. Then the eigenvalues* $\mu_1 \geq \mu_2 \geq \ldots \geq \mu_{n-1}$ *of* H *interlace with those of* G*; that is*

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \ldots \geq \mu_{n-1} \geq \lambda_n.$$

For one of our constructions we shall also make use of the Courant-Weyl Inequalities.

Theorem 4 (Courant-Weyl Inequalities, see [15], Theorem 34.2.1)**.** *For two Hermitian* $n \times n$ *matrices* A *and* B *we have*

$$\begin{array}{rcll} \lambda_i(A+B) & \leq & \lambda_{i-j+1}(A) + \lambda_j(B) & (i \geq j), \\ \lambda_i(A+B) & \geq & \lambda_{i-j+n}(A) + \lambda_j(B) & (i \leq j). \end{array}$$

We also need the following definitions from graph theory: see [7] or [9]. Let G be an n-vertex graph or multigraph. The *line graph* $L(G)$ is the graph whose vertices are the edges of G, with two vertices in $L(G)$ adjacent whenever the corresponding edges in G have exactly one vertex in common. The *root graph* of a line (multi)graph $L(G)$ is simply G itself (with a small number of exceptions, G is uniquely determined by $L(G)$). Now let $a_1, \ldots, a_n$ be non-negative integers. The *generalized line graph* $L(G; a_1, \ldots, a_n)$ is the graph $L(\hat{G})$, where $\hat{G}$ is the multigraph $G(a_1, \ldots, a_n)$ obtained from G by adding a_i pendant 2-cycles at vertex v_i $(i = 1, \ldots, n)$. An *internal path* of a graph G is a sequence of vertices $v_1, \ldots, v_k$ of G such that all vertices are distinct (except possibly v_1 and v_k), v_i is adjacent to v_{i+1} (for $i = 1, \ldots, k-1$), v_1 and v_k have degree at least 3, and all of $v_2, \ldots, v_{k-1}$ have degree 2. The following lemma collects some important results on eigenvalues that we will need.

Lemma 5. *(i)* (Perron-Frobenius, see [9], Theorem 8.8.1) *Let* H *be a proper subgraph of a connected graph* G*. Then* $\lambda_1(G) > \lambda_1(H)$*. Also,* $\lambda_1(G) > \lambda_2(G)$*.*

(ii) (Subdivision, see [11], Proposition 2.4) *Let* H *be the graph obtained from a connected graph* G *by subdividing an edge on an internal path. If* $G \neq \tilde{D}_n$*, then* $\lambda_1(H) < \lambda_1(G)$*.*

(iii) ([1], Theorem 8.2.5(v)) *Let* $\delta(G)$ *be the minimum of the degrees of the vertices of* G *and let* $\Delta(G)$ *be their maximum. Then* $\delta(G) \leq \lambda_1(G) \leq \Delta(G)$*.*

(iv) ([9], Lemma 8.6.2) *Line graphs and generalized line graphs have all their eigenvalues in the interval* $[-2, \infty)$*.*

We define a *generalized cocktail party graph (GCP)* as a graph isomorphic to a complete graph (or clique) but with some independent edges removed; that is a graph where all vertices have degree $n-1$ or $n-2$. We will use $GCP\,(n,m)$ to denote the GCP on n vertices with m edges removed, where $0 \le m \le \lfloor n/2 \rfloor$. In $GCP\,(n,m)$, we will refer to a vertex of degree $n-1$ as being "of maximal degree".

Theorem 6 (see [7], Theorem 2.3.1 and Theorem 2.1.1)**.** *A connected graph is a generalized line graph if and only if its edges can be partitioned into GCPs such that*

(i) two GCPs have at most one common vertex;
(ii) each vertex is no more than two GCPs;
(iii) if two GCPs have a common vertex, then it is of maximal degree in both of them.

Also, a graph is a line graph if and only if its edges can be partitioned in such a way that every edge is in a clique and no vertex is in more than two cliques.

A further important tool in the proof of Theorem 12 is the following:

Proposition 7 ([13], Proposition 3.2)**.** *Let G be a connected graph with $\lambda_1 > 2$ and $\lambda_2 \le 2$, then the vertices V of G can be partitioned as $V = M \cup A \cup H$ where*

(i) the induced subgraph $G|_M$ is minimal subject to $\lambda_1(G|_M) > 2$;
(ii) the set A consists of all vertices of $G|_{A\cup H}$ adjacent to some vertex of M;
(iii) the induced subgraph $G|_H$ is cyclotomic.

3. Remarks on m-Salem graphs

If an m-Salem graph G is a trivial Salem graph (see Definition 2), then naturally we refer to G as a trivial m-Salem graph. We make the simple remark that the complete graph K_{m+3} is a trivial m-Salem graph (for each $m \ge 1$): the eigenvalues of K_{m+3} are $m+2^{(1)}$ and $-1^{(m+2)}$, where the superscript denotes the multiplicity of the eigenvalue, and deleting any m vertices leaves K_3, which is cyclotomic. We now establish the more interesting result (Proposition 8) that there are infinitely many *non-trivial* m-Salem graphs for each m. To justify the construction of these non-trivial examples we use the Courant-Weyl Inequalities (Theorem 4).

When we talk about "attaching paths" below it is meant in the following way: take a graph G with a distinguished vertex v, then attaching an a-vertex path to v results in a graph with $|V\,(G)| + a$ vertices and an edge going between v and one of the end vertices of the path.

Proposition 8. *Take K_n with $n \ge 3$ and a subset of its vertices $\{v_1, \ldots, v_s\}$ where $1 < s \le n$. To each of these vertices attach a pendant path of any finite positive length, then this graph G is a Salem graph. Furthermore it is always non-trivial and, if $s = n$, $n-1$ or $n-2$ the graph is $(n-2)$-Salem, else it is $(n-3)$-Salem.*

We can interpret this construction as being a rooted product (see [8]): take K_n to be our labelled graph on n vertices and our rooted graphs to be the s paths we are attaching along with $n-s$ copies of K_1.

Proof. The graph G has $n+m$ vertices, where m is the total number of vertices attached in the s paths. Let A be the adjacency matrix of $K_n \cup mK_1$, viewed as G with all the edges of the s paths removed, so that its eigenvalues in descending order are

$$n-1^{(1)}, 0^{(m)}, -1^{(n-1)}.$$

Note that $\lambda_2(A) = 0$. Now let B be the adjacency matrix of the union of the s paths, along with $n - s$ isolated vertices, viewed as G with all the edges of the K_n removed. We now have that $A + B$ is the adjacency matrix of G. The eigenvalues of B are all in the interval $[-2, 2]$ as B is the disjoint union of cyclotomic graphs, so $\lambda_1(B) \leq 2$.

Using the first of the inequalities in Theorem 4 and setting $i = 2$, $j = 1$ we get

$$\lambda_2(A + B) \leq \lambda_2(A) + \lambda_1(B) \leq 0 + 2 = 2\,.$$

Thus G has at most one eigenvalue greater than 2. For $n \geq 3$ and $s > 1$, the graph G is not cyclotomic. Therefore $\lambda_1(G) > 2$, and this is the only eigenvalue greater than 2.

Covering each edge of each of the s paths by a K_2, and covering the edges of the K_n part of G with the single clique K_n, we see that the edges of G can be partitioned into cliques such that no vertex is in more than two cliques. By Theorem 6 G is a line graph and so by Lemma 5(iv) the least eigenvalue is greater than or equal to -2. Therefore G is a (non-bipartite) Salem graph.

To see that the Salem graph is never trivial we firstly note that $\lambda_1(G) > \lambda_1(K_n) = n - 1$ by Lemma 5(i). We also have that $\lambda_1(G) \leq \Delta(G) = n$ by Lemma 5(iii). This result is true independent of the length of our paths. Let G' be G with a single vertex added to the end of one of the s pendant paths. Then G is a proper subgraph of G', so $\lambda_1(G) < \lambda_1(G') \leq n$. The graph G therefore always has its index bounded strictly between two consecutive integers, so can never be trivial.

To see how many vertices we need to remove to induce a cyclotomic graph, we note that K_n is not cyclotomic for $n \geq 4$, so we need to remove at least $n - 3$ of the vertices from the K_n to have any chance of producing a cyclotomic subgraph. If $s \leq n - 3$ then after removing $n - 3$ vertices that include all the vertices to which the paths were attached, we are left with a K_3 (and some shorter paths), which is indeed a cyclotomic subgraph, showing that G is $(n - 3)$-Salem. If $s = n$, $n - 1$ or $n - 2$ then removing $n - 3$ vertices is not enough, since K_3 with a path attached is not cyclotomic; but in this case after removing $n - 2$ of the vertices we are left with a union of paths and deduce that G is $(n - 2)$-Salem. □

The bounds on λ_1 in the proof show that for any integer $n \geq 2$ there are infinitely many non-bipartite Salem graphs G with $\lambda_1(G) \in (n, n + 1)$. For bipartite graphs the analogous result is implied by [12, Theorem 1]. The construction here can be made even more general. We may add edges between the loose ends of any pair of pendant paths without changing the result. We can also attach copies of K_3 instead of paths (since these are also cyclotomic), although by doing this we cannot use the same triviality argument, as $\Delta(G) = n + 1$. Moreover, the regular cocktail party graph on $2n$ vertices has eigenvalues $2n - 2^{(1)}$, $0^{(n)}$ and $-2^{(n-1)}$ so we can create similar graphs that are generalized line graphs provided we are careful to make sure that we comply with the edge partitioning in Theorem 6.

The following result is an easy corollary of [13, Theorem 7.2], using the new language of m-Salem graphs.

Theorem 9 ([13], Theorem 7.2). *All Salem trees are either* 1*-Salem or* 2*-Salem.*

We also have the following trivial consequence of interlacing.

Lemma 10. *A connected subgraph of an m-Salem graph is either cyclotomic or is m'-Salem for some $m' \leq m$. For every m-Salem graph G, there is a chain of induced subgraphs $G_1 \subseteq G_2 \subseteq \cdots \subseteq G_m = G$ such that G_i is i-Salem.*

This second sentence of this lemma gives rise to a naive growing algorithm for constructing all small m-Salem graphs.

Another observation is that all 1-Salem graphs will be planar (although those in Theorem 12 are not drawn in a such a way), but m-Salem graphs for $m \geq 2$ may not be. If a graph contains an induced subgraph topologically equivalent to either K_5 or $K_{3,3}$ then it is not planar (see [10], Theorem 11.13). Removing one vertex from a graph topologically equivalent to a K_5 leaves a graph with at least four vertices of degree 3 while the cyclotomic graphs we hope to induce have at most two. Removing a vertex from a graph topologically equivalent to a $K_{3,3}$ leaves a graph with more than one cycle, which again cannot be cyclotomic.

Having got 1-Salem graphs under control, it is natural to ask about 2-Salem graphs, and so on. A stumbling-block to progress here is that not every connected graph obtained by adding a vertex to a 1-Salem graph is Salem.

4. All bipartite 1-Salem graphs

We now begin our proof of Theorem 1, starting with part (i); the bipartite 1-Salem graphs.

Theorem 11. *Let $H_1, \ldots, H_s$ be a finite set of connected cyclotomic bipartite graphs (so excluding odd cycles from the connected cyclotomic graphs described in Lemma 2), and for each i $(1 \leq i \leq s)$ let S_i be a non-empty subset of the vertices of H_i such that all the vertices of S_i fall in the same subset of the bipartition of H_i. Form the graph G by taking the union of all the H_i along with a new vertex v, and with edges joining v to each vertex in each S_i. Then*

(i) unless G is cyclotomic, it is 1-Salem;
(ii) all connected bipartite 1-Salem graphs arise in this way.

Proof. The second part is clear: if G is a bipartite 1-Salem graph, then there exists a vertex v whose deletion leaves a cyclotomic induced subgraph, and the components of this give the H_i. The first part is a consequence of interlacing: G has at most one eigenvalue greater than 2, and being bipartite we are done. □

One can be more explicit and describe precisely which combinations of H_i and S_i result in G being cyclotomic, simply by running through the possibilities for cyclotomic G given by Lemma 2. Table 1 below shows explicitly which cyclotomic graphs can arise for each choice of H_i $(1 \leq i \leq s)$. The subsets S_i that give the graphs G are easy to spot. As the degree of v is at least s, we see that no cyclotomic graphs arise when $s \geq 5$.

5. All 1-Salem generalised line graphs

Theorem 12. *The 1-Salem generalized line graphs are precisely the graphs in Figures 2, 3 and 4.*

A quick notational point: in Figures 2–4, a dashed edge indicates a path between the endpoints of the dashed edge, having an arbitrary number of edges (perhaps even none, or perhaps with a lower bound shown); the parameter attached to a

s	$H_1, \ldots, H_s$	G	s	$H_1, \ldots, H_s$	G
1	$H_1 = E_6$	$\tilde{E}_6$	2	$H_1 = K_1$, $H_2 = P_5$	$\tilde{E}_6$
	$H_1 = P_7$	$\tilde{E}_7$		$H_1 = K_1$, $H_2 = D_6$	$\tilde{E}_7$
	$H_1 = E_7$	$\tilde{E}_7$		$H_1 = K_2$, $H_2 = P_5$	$\tilde{E}_7$
	$H_1 = P_8$	$\tilde{E}_8$		$H_1 = K_1$, $H_2 = P_7$	$\tilde{E}_8$
	$H_1 = D_8$	$\tilde{E}_8$		$H_1 = H_2 = P_4$	$\tilde{E}_8$
	$H_1 = E_8$	$\tilde{E}_8$		$H_1 = P_3$, $H_2 = D_5$	$\tilde{E}_8$
	$H_1 = P_n$	$\tilde{A}_n$		$H_1 = P_2$, $H_2 = E_6$	$\tilde{E}_8$
	$H_1 = D_n$	$\tilde{D}_n$		$H_1 = K_1$, $H_2 = E_7$	$\tilde{E}_8$
				$H_1 = D_{n_1}$, $H_2 = D_{n_2}$	$\tilde{D}_n$
3	$H_1 = H_2 = H_3 = K_2$	$\tilde{E}_6$	4	$H_1 = H_2 = H_3 = H_4 = K_1$	$\tilde{D}_4$
	$H_1 = K_1$, $H_2 = H_3 = P_3$	$\tilde{E}_7$			
	$H_1 = K_1$, $H_2 = K_2$, $H_3 = P_5$	$\tilde{E}_8$			
	$H_1 = H_2 = K_2$, $H_3 = D_{n-2}$	$\tilde{D}_n$			

TABLE 1. The cyclotomic graphs that may arise in Theorem 11, arranged by the number of components s. See [7] for the (standard) definitions of E_6, E_7, E_8, A_n, D_n; P_n is the path on n vertices.

dashed edge gives the number of edges on this path. Dotted edges and vertices are used to indicate edges and vertices that may or may not be there. Here the dotted edges and vertices always form a "snake's tongue" shape, and later on we will use the notation $\hat{a}$ to indicate a path of length a (i.e., having a edges) with two extra vertices in the shape of a snake's tongue on the loose end. Thus, for example, $G_{10}(1, 1)$ and $G_{10}(1, \hat{1})$ are shown in Figure 5. The 1-Salem generalised line graphs of Theorem 12 are presented as 25 infinite families and six sporadic graphs, but there are in fact 60 non-isomorphic infinite families when we consider all the possible options of paths with snake's tongues attached.

Also, as a simple corollary to Theorem 12 we can easily note which of the graphs in Figures 2–4 are line graphs (rather than generalised line graphs) based on the characterization in Theorem 6. These are the graphs where the edges can be partitioned into cliques rather than GCPs and there are 12 infinite families and 3 sporadic graphs.

The following trivial extension of [13, Theorem 3.4] can be used to show that all the graphs in Figures 2–4 are Salem graphs.

Lemma 13. *Suppose that G is a non-cyclotomic non-bipartite graph containing a vertex v such that the induced subgraph on $V(G) \setminus \{v\}$ is cyclotomic. Also suppose that G is in the family of graphs with least eigenvalue greater than -2, then G is a Salem graph.*

To apply Lemma 13 to the graphs in Figures 2–4, we note that all the graphs in that figure are generalised line graphs (using the characterisation in Theorem 6, for example), and in each case one readily spots a vertex v whose deletion leaves a cyclotomic induced subgraph.



$G_1\,(a,b,c)$, $a \geq 1$ or $a = \hat{0}$ $G_2\,(a,b)$, $b \geq 2$ $G_3\,(a,b,c,d)$ $G_4\,(a,b,c)\,, c \geq 1$



$G_5\,(a,b)\,,\ a,b \geq 1$ $G_6\,(a,b,c)$, $b \geq 2$ $G_7\,(a,b)$, $a,b \geq 2$ $G_8\,(a,b)$



$G_9\,(a)$, $a \geq 2$ $G_{10}\,(a,b)$ $G_{11}\,(a)$, $a \geq 2$ $G_{12}\,(a,b,c)$

FIGURE 2. Twelve of the 25 infinite families of 1-Salem generalised line graphs. The parameters a, b, c, d are ≥ 0 unless specified.

We now prove Theorem 12 using the structure given in Proposition 7; that is we will grow our graphs starting with the vertices in M, then $M \cup A$, then $M \cup A \cup H$. We quickly note a simple lemma that will be referred to frequently in the proof.

Lemma 14. *A* 1*-Salem graph* G *may not contain an induced* K_5*. Also, if it contains an induced* K_4*, then only one of the four vertices in that* K_4 *may be attached to any other vertices in* G *and the vertex we remove to make* G *cyclotomic must be this distinguished vertex in* K_4*.*

Proof. The first sentence is clear since if we remove any one of the vertices of K_5 we obtain a K_4 which is not cyclotomic. For the second sentence, removing a vertex v of K_4 leaves a K_3 so if any of the other vertices of the K_4 were attached to any vertices of G, so would the K_3 be after removing v, and no connected supergraph of K_3 is cyclotomic. ☐

5.1. M — **the minimal graphs.** In this step we will need three of the 31 minimal forbidden subgraphs for the property of being a generalized line graph (see [7, Theorem 2.3.18]). The three that we will use are in Figure 6 below.

Proposition 15. *All Salem generalized line graphs must contain a* K_3*.*



$G_{13}\,(a,b)$, $b \geq 2$ $G_{14}\,(a,b)$, $a \geq 1$ $G_{15}\,(a)$

$G_{16}\,(a,b)$ $G_{17}\,(a)$, $a \geq 1$ $G_{18}\,(a)$ $G_{19}\,(a,b)$

$G_{20}\,(a)$, $a \geq 2$ $G_{21}\,(a)$ $G_{22}\,(a)$ $G_{23}\,(a,b)$

$G_{24}\,(a)$, $a \geq 2$ $G_{25}\,(a)$

FIGURE 3. Thirteen of the 25 infinite families of 1-Salem generalised line graphs. The parameters a, b are ≥ 0 unless specified.



G_{26} G_{27} G_{28} G_{29} G_{30} G_{31}

FIGURE 4. The six sporadic 1-Salem generalised line graphs.

Proof. Lemma 5(iv) tell us that all generalized line graphs have their least eigenvalue $\lambda_n \geq -2$ and by definition Salem graphs have $\lambda_1 > 2$, so a Salem generalized line graph must be non-bipartite and therefore must contain an odd cycle. Let C be an odd cycle of shortest length in a connected Salem generalised line graph G. The subgraph of G induced by the vertices of C is just C, else there would be a shorter odd cycle in G. Since an odd cycle is cyclotomic, G must contain a vertex



FIGURE 5. An illustration of the hat convention.



FIGURE 6. Three of the minimal forbidden subgraphs for the property of being a generalized line graph. The names for these graphs are taken from [7].

v adjacent to a vertex a on C (and perhaps adjacent to other vertices on C). If C has length 3, we are done.

The next case is to consider when C has length 5. If v were adjacent only to a on C, then C together with v would induce $G^{(13)}$, one of the minimal forbidden subgraphs, so we must add at least one more edge between v and a vertex on C. If we have an edge between v and either of the neighbours of a on C, then we would form a triangle, so from now on we may suppose that v is not adjacent to either of these neighbours of a. Similarly, if we attach v to both of the vertices on C that are not adjacent to a, then we would form a triangle. But we cannot attach v to exactly one of these two non-neighbours of a on C, else we would have $G^{(1)}$ as an induced subgraph. We conclude that C cannot have length 5.

For C of length 7 or greater we follow a similar method. If v were adjacent only to a on C, then C together with v would induce a $G^{(12)}$, one of the minimal forbidden subgraphs, so we must add at least one more edge between v and a vertex on C. If v were adjacent to a vertex x on C that was neither one of the neighbours of a on C, nor one of the vertices distance two along C from a, then one could find a shorter odd cycle in G (using a, v, x, and completing the cycle along C in the appropriate direction to make the length odd), so we can exclude this possibility. If v were adjacent to a neighbour of a on C, then we would have a triangle. Since this is not the case, v must be adjacent to one (or both) of the vertices on C that are distance 2 from a, but not to either of the neighbours of a on C, giving $G^{(1)}$ as an induced subgraph. □

As a corollary to this we get that the minimal graphs we are after must be the three ways of attaching a single vertex to a K_3.

Corollary 16. *The minimal graphs with respect to the property of being a Salem generalized line graph are the three graphs in Figure 7.*

We now know what our minimal graphs look like but before proceeding we make one more observation. For $G \setminus v$ to be cyclotomic we must have $v \in M$; that is, the vertex we are removing to induce a cyclotomic graph must be one of the vertices of the minimal graph M.



FIGURE 7. The three minimal graphs in Corollary 16.

5.2. A — **the adjacent vertices.** Growing our graph from $G|_M$ to $G|_{M\cup A}$ is the most difficult part, however the restriction to 1-Salem generalized line graphs reduces this to a finite search, after some work. By Theorem 6 we have a highly-constrained structure: we cannot simply add vertices and edges anywhere. First let us consider the ways we can partition M_1, M_2 and M_3 into GCPs; this will reveal where we can add vertices. The generalised line graph M_1 can be seen uniquely as a K_3 and a K_2; it then has three vertices of maximal degree that are not already in two GCPs. And M_3 can be seen uniquely as a K_4, but by Lemma 14 we know that we can attach further vertices to only one of its original vertices. The graph M_2, however, is one of the seven graphs that has two non-isomorphic root multigraphs (see [7, Theorem 2.3.4]) so we need to consider both versions. Let $M_{2,1}$ be M_2 partitioned as a K_3 with two K_2's attached and both joined at their other ends; here we only have one vertex that is of maximal degree and not already in two GCPs. Let $M_{2,2}$ be M_2 partitioned as a $GCP\,(4,1)$ where we then have two vertices of maximal degree.

The set of vertices A are those in the graph $G|_{A\cup H}$ that are adjacent to a vertex in M. By Theorem 6, to grow from $G|_M$ to $G|_{M\cup A}$ we can:

- expand a GCP to a larger one that contains the original one (taking care of the degrees of vertices);
- attach a new GCP to a vertex of maximal degree that is only in one GCP (attaching only at vertices of maximal degree);
- do both of the above.

Once we have done this we can then also add an edge between any two vertices of A of maximal degree in their GCPs (in effect, adding a K_2) or take two maximal degree vertices of A that are in separate GCPs and only in one GCP each and "merge" them together (connect two disconnected GCPs by making them share an available maximal degree vertex).

Lemma 14 helps here as we know that we need not consider any GCPs that contain a K_5. In fact, by studying the GCPs G that have this property and looking at the induced graph $G-v$ for each $v \in V\,(G)$ we find the fairly short list of GCPs that we can attach or expand to in Table 2 below.

Note that $GCP\,(4,2) = C_4$ is not included; when partitioned as $GCP\,(4,2)$ it has no vertices of maximal degree, so cannot be attached to anything and when partitioned as four K_2's it has no vertices that are not already in two GCPs. Also, $GCP\,(3,1)$ and $GCP\,(5,2)$ are two of the seven graphs that have non-isomorphic root multi-graphs. However, when $GCP\,(5,2)$ is partitioned as two K_3's and two K_2's it has no vertices that are not already in two GCPs and when $GCP\,(3,1)$ is partitioned as two K_2's one of its vertices will not be in A.

The process of going from $G|_M$ to $G|_{M\cup A}$ is then a finite one; we only have so many GCPs in the minimal graphs M_1, $M_{2,1}$, $M_{2,2}$ and M_3 to expand and only so

$GCP\,(2,0) = K_2$		2 vertices of maximal degree
$GCP\,(3,1)$		1 vertex of maximal degree
$GCP\,(3,0) = K_3$		3 vertices of maximal degree
$GCP\,(4,1)$		2 vertices of maximal degree
$GCP\,(4,0) = K_4$		4 vertices of maximal degree
$GCP\,(5,2)$		1 vertex of maximal degree

Table 2. The GCPs that we can attach or expand to in A.

many ways to attach these six GCPs to them. We also have a small number of cases where we can add in extra edges between vertices of A or merge them. In working through all these combinations we discard a number of graphs that are not 1-Salem as they require more than one vertex to be removed to make them cyclotomic.

The list of 1-Salem generalised line graphs $G|_{M\cup A}$ has been omitted for reasons of space. There are 224 that are distinct as graphs, although many of these graphs can arise in more than one way as $G|_{M\cup A}$. The largest has 11 vertices. Initially these were all found by hand; the list was then checked by a computer search.

5.3. H — **the cyclotomic parts.** We now look at the set H in Lemma 7. We can reduce our choices by observing that the only cyclotomic graphs that are also generalized line graphs are $\tilde{D}_n$ and $\tilde{A}_n = C_{n+1}$. However, we will show that $G|_H$ cannot contain cycles and can only contain subgraphs of $\tilde{D}_n$. For $n > 3$ we note that $\tilde{A}_n$ can be partitioned uniquely as $(n+1)$ K_2's with each vertex in two GCPs, so we cannot simply attach vertices to it. The other option is then to expand a GCP to a larger one so that we now have vertices only in one GCP to attach to other things. The smallest case is to expand one of the $(n+1)$ K_2's to a K_3. Clearly, this new vertex must be in A rather than H as the graph is no longer cyclotomic. However, in order to make the graph cyclotomic we must remove a vertex but all of the vertices are in A or H and we have shown that the vertex we remove must be in M. Cycles of length 3 can be seen as both three K_2's or one K_3 but by similar reasoning on the choice of vertex we are removing, we can exclude this case too. A similar argument holds again for cycles of length 4 for both ways of partitioning its edges.

The graph $\tilde{D}_n$ can be uniquely partitioned as $(n-4)$ K_2's with a $GCP\,(2,0)$ (or snake's tongue) at either end. However, in this graph each vertex of maximal degree within its GCP is already in two GCPs. If we remove one or both of the snake's tongues we are left with graphs we can work with—a path or a path with a snake's tongue on the end—each with at least one vertex of maximal degree that is only in one GCP. With this in mind we can think of $G|_H$ simply consisting of paths of any length, possibly with a snake's tongue on one end.

The final step in growing these graphs is to go from $G|_{M\cup A}$ to $G|_{M\cup A\cup H} = G$. To any vertices in A of maximal degree and only in one GCP we can attach a single path of arbitrary length (remembering that Lemma 14 tells us that we can only attach things to one vertex of a K_4). If we so choose, we can join any two of these pendant paths together (equivalently, attach a path or arbitrary length to two different elements of A of maximal degree that are each only in one GCP). Furthermore, on the end of any pendant paths we can include a snake's tongue. These graphs are then precisely the graphs in Figures 2–4 and Lemma 13 tells us that they are all 1-Salem, completing this section of the proof.

We can consider letting the lengths of one or more pendant or internal paths tend to infinity. By [13, Corollary 4.4] the corresponding sequence of graph Salem numbers converges to a Pisot number. By studying the different ways we can let paths tend to infinity, we can produce several explicit families of Pisot numbers.

6. The 1-Salem graphs represented in E_8

6.1. **The root system and signed graph E_8.** There are various concrete descriptions of the root system E_8 (see [3, Chapter 3], for example). Here is one of them. Let $\mathbf{e}_1, \ldots, \mathbf{e}_8$ be an orthogonal basis for $\mathbb{R}^8$, with each $\mathbf{e}_i$ having length $\sqrt{2}$. The root system E_8 contains 240 vectors: the 16 vectors $\pm\mathbf{e}_1, \ldots, \pm\mathbf{e}_8$, and 224 vectors of the form $(\pm\mathbf{e}_i \pm \mathbf{e}_j \pm \mathbf{e}_k \pm \mathbf{e}_l)/2$, where $ijkl$ is one of the 14 strings 1234, 1256, 1278, 1357, 1368, 1467, 2358, 2367, 2457, 2468, 3456, 3478, 5678. The usual inner product of any two distinct vectors from these 240 is either -1, 0, or 1; and all have length 2. Hence we can define a signed graph (which we might as well also call E_8) with 240 vertices corresponding to these vectors, and with adjacencies of distinct vertices given by the inner product of the corresponding vectors.

The *Weyl group* $W(E_8)$ is the group of isometries of E_8 generated by the involutions $\mathbf{v} \mapsto \mathbf{v} - (\mathbf{v} \cdot \mathbf{v}_i)\mathbf{v}_i$ for each of the 240 vertices $\mathbf{v}_i$ (in fact there are only 120 of these involutions, since $\mathbf{v}_i$ and $-\mathbf{v}_i$ yield the same involution).

A graph is said to be represented in E_8, if it appears as an induced subgraph of the signed graph E_8. If A is the adjacency matrix of such a graph, then $A + 2I$ is the Gram matrix of some subset of the vectors forming the root system E_8.

6.2. **Fast backtracking in E_8.** Let P be a property of graphs such that if G has property P then so does any induced subgraph. (Such a property is called *hereditary*.) The following process provides a computationally-efficient method for searching for all graphs that: (a) are represented in E_8; and (b) have property P.

Assign an ordering to the vertices of E_8: $\mathbf{v}_1, \ldots, \mathbf{v}_{240}$. A naive search for all graphs having property P that are represented in E_8 would be as follows. Start with an empty list of vertices; look through the vertices in order, adding them to the list so long as the signed subgraph induced by the vertices in the list is actually a graph and has property P; once stuck, backtrack. The number of signed subgraphs of E_8 is so large that this process is only practical for extremely restrictive properties P. To improve matters, we exploit the isometries of E_8.

Choose a set S of isometries of E_8. When considering if it is possible to add a new vertex to ones list, apply each isometry in S to the potential enlarged list, and reject the new vertex if any of the permuted lists appears earlier in the lexicographical ordering derived from our ordering of the vertices. Larger sets S make each update more expensive, but potentially reduce the backtracking by pruning certain branches at an early stage.

We used a few other programming tricks to speed up the process further, but the main idea was to exploit isometries of E_8. In practice we used a set S of size 162, including the 120 involutions above, and 42 carefully selected others.



Figure 8. The two 11-vertex 1-Salem graphs that are not generalised line graphs.

6.3. Cyclotomic graphs represented in E_8.

Proposition 17. *Let G be a cyclotomic graph represented in E_8. Then G has at most 10 vertices.*

Proof. The fast backtracking approach described above was applied to find all cyclotomic graphs represented in E_8. The largest ones had 10 vertices. ☐

There are three distinct 10-vertex examples of cyclotomic graphs represented in E_8, up to isomorphism. They are: (i) two 4-cycles and two isolated vertices; (ii) two copies of $\tilde{D}_4$; (iii) a 4-cycle and $\tilde{D}_5$. The largest connected examples have 9 vertices: (i) $\tilde{D}_8$; (ii) $\tilde{E}_8$.

6.4. **1-Salem graphs that are not generalised line graphs.** By [2] we know that all of these are represented in E_8. Knowing the small size of cyclotomic graphs represented in E_8 restricts our search to graphs with at most 11 vertices. We could, in principle, modify our search for cyclotomics to deal with this harder problem. Or we could, in principle, search for all connected 1-Salem subgraphs of the 473 graphs in [7, Table A6] (the maximal connected graphs subject to having all eigenvalues at least -2). But a list of all Salem graphs of up to 12 vertices has been computed by the first author, and the simplest strategy for us was therefore to look through this list for non-bipartite 1-Salem graphs, and then see which of these were not generalised line graphs. For this latter point, the characterisation by excluded subgraphs (see [6], [16]) was used—slightly simplified by the fact that 11 of the 31 excluded subgraphs have eigenvalues below -2, so cannot appear as subgraphs of non-bipartite Salem graphs (see [7, Theorem 2.3.19]).

Table 3 shows the number of non-bipartite connected 1-Salem graphs that are not generalised line graphs by the number of vertices. The largest examples have 11 vertices; they are shown in Figure 8. These computations complete the final part of the proof of Theorem 1, which we state here as a theorem.

Theorem 18. *There are 377 non-bipartite connected 1-Salem graphs that are not generalised line graphs. The numbers of vertices for these graphs range between 6 and 11; the number of graphs for each of these numbers of vertices is shown in Table 3.*

Number of vertices	6	7	8	9	10	11
Number of graphs	10	43	111	153	58	2

Table 3. The number of non-bipartite connected 1-Salem graphs that are not generalised line graphs.

Number of vertices	6	7	8	9	10
Number of graphs	1	4	7	8	5

TABLE 4. The number of non-bipartite triangle-free Salem graphs.

7. TRIANGLE-FREE NON-BIPARTITE SALEM GRAPHS

Let G be a triangle-free non-bipartite Salem graph. Being a non-bipartite Salem graph, G is either a generalised line graph or is represented in E_8. Being a triangle-free Salem graph, G is not a generalised line graph by Proposition 15. Hence G must be represented in E_8. In particular, only finitely many such graphs exist.

By [14, Section 7.3], triangle-free subgraphs of E_8 have at most 16 vertices, and restricting to those that are not cyclotomic, the maximum number of vertices is 12. Again it was convenient for us here simply to search the first author's list of small Salem graphs. This established the following Proposition.

Proposition 19. *There are exactly* 25 *non-bipartite triangle-free Salem graphs. All have at most* 10 *vertices. The numbers of each size are given in Table 4.*

The 10-vertex examples include the Petersen graph: this is the only case that is a trivial Salem graph (see Definition 2; moreover it is an integral graph). All but one of the 25 graphs are planar.

8. THE TRIVIAL NON-BIPARTITE 1-SALEM GRAPHS

For the non-bipartite graphs it is possible to observe when they are trivial Salem numbers (when $\lambda_1 \in \mathbb{Z}$). This is a different question to that of integral graphs as we only require the largest eigenvalue to be an integer, however the two problems intersect. For the finite number of graphs from E_8 and the sporadic generalized line graphs we simply need to calculate the eigenvalues to see when they are trivial. For the infinite families however we need to take a different approach. Similarly to the proof of Proposition 8, we look to bound the largest eigenvalue between two consecutive integers when possible. It turns out we can always do this for the 25 infinite families of 1-Salem generalized line graphs.

Lower bounds are easily found by looking at the graphs when all pendant paths and cycles are at their smallest. Upper bounds are found by calculating the largest eigenvalue of an "extremal" version of the graph G_E; that is, one where all the pendant paths are infinitely long and all internal paths (cycles and paths with snake's tongues) are as short as possible. We then use Lemma 5: subdivision of internal paths tells us that $\lambda_1(G) < \lambda_1(G_E)$ as more vertices are added to the path, and Perron-Frobenius tells us that $\lambda_1(G) < \lambda_1(G_E)$ for pendant paths of any finite length. Subdivision followed by Perron-Frobenius deals with graphs that contain both internal paths and pendant paths. To calculate the largest eigenvalue of the graphs G_E that are infinite we use the following result from [13, Lemma 4.1] on reciprocal polynomials

$$R_{G_p}(z) = \frac{z^{2p}-1}{z^2-1} R_{G_1}(z) - \frac{z^{2p}-z^2}{z^2-1} R_{G_0}(z),$$

where $R_G(z) = z^n \chi_G(z+1/z)$ and G_p denotes the graph G with a pendant path of length p attached. We apply this as many times as necessary and let the lengths of the paths tend to infinity. Our $\lambda_1(G_E) = z_1 + 1/z_1$ where z_1 is the largest root of $R_{G_E}(z)$. The trivial graphs can be seen in Figure 9.



$G_3\left(\hat{0},\hat{0},\hat{0},\hat{0}\right)$ $G_4\left(\hat{0},\hat{0},2\right)$ $G_5\,(2,2)$ $G_6\left(\hat{0},2,\hat{0}\right)$ $G_7\,(2,2)$ $G_{10}\left(\hat{0},\hat{0}\right)$

$G_{11}\,(2)$ $G_{18}\,(0)$ G_{31} E_1 E_2 E_3 E_4

E_5 E_6 E_7

Figure 9. The trivial non-bipartite 1-Salem graphs

With the exceptions of the graphs in Figure 9 we can bound the indices of graphs G_1, G_2, G_3, $G_6, \ldots, G_{11}$ and G_{15} strictly between 2 and 3 and graphs G_{13}, G_{14}, $G_{16}, \ldots, G_{25}$ strictly between 3 and 4. The graphs G_4, G_5 and G_{12} have indices between 2 and 4 but the only times they give trivial graphs are the two in Figure 9.

It is worth noting that all but five of these trivial Salem graphs are in fact integral. The graphs $G_4(\hat{0},\hat{0},2)$, E_2, E_4, E_5 and E_7 are non-integral. With the exception of G_{31}, the remaining 10 graphs overlap with those found in [5] that are also 1-Salem. Any integral graph with index 3 which also has the property that on removing one vertex it becomes cyclotomic must be 1-Salem by Lemma 5(i). The graph G_{31} has index 4 but we will not be able to apply the same comment to a similar list as 3 or -3 may be eigenvalues too.

References

[1] B. Bollobás, *Graph theory: an introductory course*, Graduate Texts in Mathematics **63**, Springer, New York, 1979.

[2] P.J. Cameron, J.M. Goethals, J.J. Seidel, E.E. Shult, Line graphs, root systems, and elliptic geometry, *J. Algebra* **43** (1976), 305–327.

[3] P.J. Cameron, J.H. van Lint, *Designs, graphs, codes and their links*, Cambridge University Press, Cambridge, 1991.

[4] A.L. Cauchy, Sur l'équation à l'aide de laquelle on détermine les inégalitiés séculaires des mouvements des planètes, Oeuvres complètes, Iièeme Série, **9**, 174–195, Gauthier-Villars, 1829.

[5] T. Chung, J. Koolen, Y. Sano, T. Taniguchi, The non-bipartite integral graphs with spectral radius three, *Linear Algebra and its Applications* **435**(10) (2011), 2544–2559.

[6] D. Cvetković, M. Doob, S. Simić, Generalised line graphs, *J. Graph Theory* **5**, 385–399, 1981.

[7] D. Cvetković, P. Rowlinson, S. Simić, *Spectral generalizations of line graphs: on graphs with least eigenvalue* -2, London Mathematical Society Lecture Note Series **314**, Cambridge University Press, Cambridge, 2004.

[8] C. Godsil, B. McKay, A new graph product and its spectrum, *Bull. Australian Math. Soc.* **18**(01) (1978), 21–28.

[9] C. Godsil, G. Royle, *Algebraic Graph Theory*, Springer, New York, 2001.

[10] F. Harary, *Graph Theory*, Addison-Wesley, 1969.

[11] A.J. Hoffman, J.H. Smith, On the spectral radii of topologically equivalent graphs, in *Recent advances in graph theory*, 273–281, ed. M. Fiedler, Academia Praha, 1975.

[12] J. McKee, C. Smyth, P. Rowlinson, Salem numbers and Pisot numbers from stars, in *Number Theory in Progress*, vol. 1 (Zakopane-Kocielisko, 1997), edited by K. Győry, H. Iwaniec, J. Urbanowicz, 309–319, de Gruyter, Berlin, 1999.

[13] J. McKee, C. Smyth, Salem numbers, Pisot numbers, Mahler measure, and graphs, *Experimental Mathematics* **14** (2005), 211–229.

[14] J. McKee, C. Smyth, Integer symmetric matrices having all their eigenvalues in the interval $[-2, 2]$, *J. Algebra* **317** (2007), 260–290.

[15] V. Prasolov, S. Ivanov, *Problems and theorems in linear algebra*, American Mathematical Society, 1994.

[16] S.B. Rao, N.M. Vinghi, K.S. Vijayan, The minimal forbidden subgraphs for generalized line-graphs, in *Combinatorics and Graph Theory, Proc. Symp. Indian Statistical Institute, Calcutta, 25–29 February 1980*, edited by S.B. Rao, 459–472, Lecture Notes in Math. **855**, Springer, Berlin, 1981.

[17] J.H. Smith, Some properties of the spectrum of a graph, in *Combinatorial structures and their applications (Proc. Calgary Internat. Conf., Calgary, Alta., 1969)*, edited by R. Guy, H. Hanani, H. Saver, J. Schönheim, 403–406, Gordon and Breach, New York, 1970.

Department of Mathematics, Royal Holloway, University of London, Egham Hill, Egham, Surrey, TW20 0EX, England, UK.
E-mail address: `Jonathan.Cooley.2009@rhul.ac.uk`

Department of Mathematics, Royal Holloway, University of London, Egham Hill, Egham, Surrey, TW20 0EX, England, UK.
E-mail address: `Lee.Gumbrell.2009@rhul.ac.uk`

Department of Mathematics, Royal Holloway, University of London, Egham Hill, Egham, Surrey, TW20 0EX, England, UK.
E-mail address: `James.McKee@rhul.ac.uk`